



TITLE:

Synthesis of cross-conjugated trienes by rhodium-catalyzed dimerization of monosubstituted allenes

AUTHOR(S):

Miura, Tomoya; Biyajima, Tsuneaki; Toyoshima, Takeharu; Murakami, Masahiro

CITATION:

Miura, Tomoya ...[et al]. Synthesis of cross-conjugated trienes by rhodium-catalyzed dimerization of monosubstituted allenes. BEILSTEIN JOURNAL OF ORGANIC CHEMISTRY 2011, 7: 978-981

ISSUE DATE:

2011-05

URL:

<http://hdl.handle.net/2433/163438>

RIGHT:

© 2011 Miura et al; licensee Beilstein-Institut. This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The license is subject to the Beilstein Journal of Organic Chemistry terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

Synthesis of cross-conjugated trienes by rhodium-catalyzed dimerization of monosubstituted allenes

Tomoya Miura, Tsuneaki Biyajima, Takeharu Toyoshima
and Masahiro Murakami*

Full Research Paper

Open Access

Address:
Department of Synthetic Chemistry and Biological Chemistry, Kyoto
University, Katsura, Kyoto 615-8510, Japan

Email:
Tomoya Miura - tmiura@sbchem.kyoto-u.ac.jp; Masahiro Murakami* -
murakami@sbchem.kyoto-u.ac.jp

* Corresponding author

Keywords:
allene; cross-conjugated triene; dimerization; rhodium; stereoselective

Beilstein J. Org. Chem. **2011**, *7*, 578–581.
doi:10.3762/bjoc.7.67

Received: 25 January 2011
Accepted: 30 March 2011
Published: 09 May 2011

This article is part of the Thematic Series "Allene chemistry".

Guest Editor: K. M. Brummond

© 2011 Miura et al; licensee Beilstein-Institut.
License and terms: see end of document.

Abstract

A rhodium(I)/dppe catalyst promoted dimerization of monosubstituted allenes in a stereoselective manner to give cross-conjugated trienes, which are different from those obtained by a palladium catalyst.


Introduction

Cross-conjugated trienes, known as [3]dendralenes [1], are attractive synthetic precursors used for consecutive double [4 + 2] cycloaddition reactions [2–4] to provide a rapid access to polycyclic carbon frameworks. Thus, a number of methods for the preparation of the parent 3-methylenepenta-1,4-diene [5] and its substituted derivatives [6–17] has been developed. Among these, transition-metal-catalyzed dimerization of allenes presents a unique entry to substituted cross-conjugated trienes. For example, a nickel(0)/triphenylphosphine complex catalyzes a dimerization reaction of 3-methylbuta-1,2-diene to afford 2,5-dimethyl-3,4-bismethylenehex-1-ene [18,19]. The nickel-catalyzed reaction, however, leads to a complex mixture of products when monosubstituted allenes such as penta-1,2-diene and 1-phenylpropa-1,2-diene are employed [20]. On the other

hand, a palladium-catalyzed dimerization reaction of monosubstituted allenes produces substituted cross-conjugated trienes **2** in high yield (Scheme 1) [21]. We report here that dimerization of monosubstituted allenes is also catalyzed by a rhodium(I)/dppe complex to form cross-conjugated trienes **3**, which are different from those obtained with the palladium catalyst.


Results and Discussion

We initiated our study using undeca-1,2-diene (**1a**) as the model substrate and a rhodium(I) complex as the catalyst (Table 1). When **1a** was treated with a catalytic amount of [RhCl(cod)]₂ (2.5 mol %, cod = cycloocta-1,5-diene) in toluene at 130 °C for 12 h, **2a** was formed in 40% NMR yield along with another minor dimerized product (13% NMR yield) and unidentified



compounds (Table 1, entry 1). The structure of the minor dimerized product was determined to be (*E*)-10,11-dimethylenecos-8-ene (**3a**) by ^1H and ^{13}C NMR spectroscopy. Thus, the two isomeric dimers, one identical to the isomer obtained by the palladium-catalyzed reaction and the other a different isomer, were produced by the rhodium-catalyzed reaction. Next, several phosphine ligands were examined (Table 1, entries 2–5). To our delight, the use of the dppe ligand suppressed the formation of **2a** and the unidentified compounds, and increased the NMR yield of **3a** to 96% (86% isolated yield, Table 1, entry 4). A complex mixture of products was obtained when the reaction temperature was lowered from 130 °C to 90 °C (Table 1, entry 6). Moreover, the use of $[\text{Rh}(\text{OH})(\text{cod})]_2$ and $\text{Rh}(\text{acac})(\text{cod})$ as the precatalyst resulted in a decrease of the reaction rate (Table 1, entries 7 and 8).

Table 1: Optimization of reaction conditions^a.



Entry	X	Ligand ^b	T (°C)	Yield of 2a (%) ^c	Yield of 3a (%) ^c
1	Cl	none	130	40	13
2	Cl	PPh_3^d	130	24	18
3	Cl	dppe	130	24	37
4	Cl	dppe	130	<5	96 (86)
5	Cl	dppp	130	17	50
6	Cl	dppe	90	38	24
7	OH	dppe	130	40	10
8	acac ^e	dppe	130	44	<5

^aReactions conducted on a 0.4 mmol scale.


^bdppe = 1,1-bis(diphenylphosphino)methane, dppe = 1,2-bis(diphenylphosphino)ethane, dppp = 1,3-bis(diphenylphosphino)propane.

^cNMR yield using mesitylene as an internal standard. Isolated yield given in parenthesis.

^dUsing 10 mol % of PPh_3 .

^eUsing 5.0 mol % of $\text{Rh}(\text{acac})(\text{cod})$.

We propose that the dimerization reaction proceeds through the pathway outlined in Scheme 2. Initially, two molecules of **1a** coordinate to a rhodium(I) center at the terminal carbon–carbon double bonds from their sterically less-hindered sides. Oxidative cyclization occurs in a head-to-head manner to form the five-membered rhodacyclic intermediate **A** [22–25], which is in equilibrium with another rhodacyclic intermediate **B** via σ – π – σ isomerization. Then, β -hydride elimination takes place with **B** to form rhodium hydride **C** stereoselectively. Finally, reductive elimination from **C** yields **3a** together with the catalytically active rhodium(I) complex. It is also conceivable, however, that oxidative cyclization of two molecules of **1a** occurs in a tail-to-tail manner to directly furnish **B**. The other isomer **2a** could be formed through allylic 1,3-migration of rhodium from **C** and subsequent reductive elimination.



Scheme 2: A proposed reaction pathway.

Under the optimized reaction conditions using dppe as the ligand, various monosubstituted allenes **1b–j** were subjected to the catalytic dimerization reaction (Table 2). In most cases, essentially one isomer **3** was formed, and the other isomer **2** was barely detectable in the ^1H NMR spectrum of the crude reaction mixture (<5%). Allenes **1b–i** possessing a primary alkyl group reacted well to afford the corresponding products **3b–i** in yields ranging from 63% to 90% (Table 2, entries 1–8). Functional groups such as benzyloxy, siloxy, hydroxy and cyano groups were tolerated in the alkyl chain under the reaction conditions. Cyclohexylpropa-1,2-diene (**1j**) possessing a secondary alkyl group also participated in the dimerization reaction (Table 2, entry 9). On the other hand, 1,1-disubstituted allenes such as 3-methylbuta-1,2-diene and 3-pentylcta-1,2-diene failed to undergo the dimerization reaction, in contrast to the nickel-catalyzed reaction [18,19].

Table 2: Synthesis of cross-conjugated trienes by the allene dimerization reaction^a.

Entry	1	R	R'	3	Yield (%) ^b
1	1b	C ₅ H ₁₁	H	3b	85
2	1c	CH ₂ Ph	H	3c	78 ^c
3	1d	Cy	H	3d	83 ^d
4	1e	CH ₂ OBn	H	3e	70
5	1f	(CH ₂) ₃ OBn	H	3f	78
6	1g	(CH ₂) ₃ OSiMe ₂ <i>t</i> -Bu	H	3g	90
7	1h	(CH ₂) ₃ OH	H	3h	63
8	1i	(CH ₂) ₃ CN	H	3i	75
9	1j	–(CH ₂) ₅ –		3j	60 ^d

^aReactions conducted on a 0.4 mmol scale.

^bIsolated yield unless otherwise noted.

^cThe product was accompanied by a small amount of an unidentified impurity.

^dNMR yield using mesitylene as an internal standard.

Next, we examined the consecutive double [4 + 2] cycloaddition reaction of the cross-conjugated trienes obtained in the present study. Triene **3a** was treated with 4-phenyl-1,2,4-triazoline-3,5-dione (**4**, PTAD), a highly reactive dienophile, in toluene at 0 °C (Scheme 3). The conversion of **3a** was complete within 1 h, and after chromatographic isolation, bisadducts **5a**

and **5a'** were obtained in 75% and 6% yields, respectively. The major bisadduct **5a** resulted from initial addition to the more congested diene moiety of **3a** (site β). When tetracyanoethylene (**6**, TCNE), which was a less reactive dienophile than **4**, was used, [4 + 2] cycloaddition also occurred preferentially at site β, but only once on heating at 60 °C for 24 h.


Conclusion

In summary, we have developed a new dimerization reaction of monosubstituted allenes catalyzed by a rhodium(I)/dppe complex, allowing the stereoselective formation of substituted cross-conjugated trienes. It is interesting that the rhodium catalyst and the palladium catalyst gave different types of cross-conjugated trienes.


Experimental

General procedure for rhodium-catalyzed dimerization of monosubstituted allenes

To a side-arm tube equipped with a stirrer bar, was added [RhCl(cod)]₂ (4.9 mg, 2.5 mol %) and dppe (7.7 mg, 5 mol %). The tube was evacuated and refilled with argon three times. Then, toluene (4 mL) and substrate **1** (0.4 mmol) were added via syringe and the tube was closed. After heating at 130 °C for 6 h, the reaction mixture was cooled to room temperature, passed through a pad of Florisil[®] and eluted with ethyl acetate (≈ 90–100 mL). The filtrate was concentrated under reduced pressure and the residue purified by preparative thin-layer chromatography to give product **3**. Although the isolated **3** was relatively labile, it could be kept at –30 °C for days without any detectable decomposition or polymerization.



Scheme 3: [4 + 2] cycloaddition reaction of **3a** with PTAD and TCNE.



Supporting Information

Supporting Information File 1

Experimental details and spectroscopic data for new compounds.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-7-67-S1.pdf>]

Acknowledgements

This work was supported in part by MEXT (Grant-in-Aid for Challenging Exploratory Research, No. 21656033). T. T. is grateful for Research Fellowship from JSPS for Young Scientists.

References

- Hopf, H. *Nature* **2009**, *460*, 183–184. doi:10.1038/460183a
- Tsuge, O.; Wada, E.; Kanemasa, S. *Chem. Lett.* **1983**, *12*, 239–242. doi:10.1246/cl.1983.239
- Kwon, O.; Park, S. B.; Schreiber, S. L. *J. Am. Chem. Soc.* **2002**, *124*, 13402–13404. doi:10.1021/ja028086e
- Brummond, K. M.; You, L. *Tetrahedron* **2005**, *61*, 6180–6185. doi:10.1016/j.tet.2005.03.141
- Bradford, T. A.; Payne, A. D.; Willis, A. C.; Paddon-Row, M. N.; Sherburn, M. S. *J. Org. Chem.* **2010**, *75*, 491–494. doi:10.1021/jo9024557
- Woo, S.; Squires, N.; Fallis, A. G. *Org. Lett.* **1999**, *1*, 573–576. doi:10.1021/ol990695c
- Chang, H.-M.; Cheng, C.-H. *J. Org. Chem.* **2000**, *65*, 1767–1773. doi:10.1021/jo991651k
- Brummond, K. M.; Chen, H.; Sill, P.; You, L. *J. Am. Chem. Soc.* **2002**, *124*, 15186–15187. doi:10.1021/ja027588p
- Chin, C. S.; Lee, H.; Park, H.; Kim, M. *Organometallics* **2002**, *21*, 3889–3896. doi:10.1021/om020410r
- Kang, B.; Kim, D.-H.; Do, Y.; Chang, S. *Org. Lett.* **2003**, *5*, 3041–3043. doi:10.1021/ol035014z
- Shi, M.; Shao, L.-X. *Synlett* **2004**, 807–810. doi:10.1055/s-2004-820017
- Bräse, S.; Wortal, H.; Frank, D.; Vidović, D.; de Meijere, A. *Eur. J. Org. Chem.* **2005**, 4167–4178. doi:10.1002/ejoc.200500330
- Bradford, T. A.; Payne, A. D.; Willis, A. C.; Paddon-Row, M. N.; Sherburn, M. S. *Org. Lett.* **2007**, *9*, 4861–4864. doi:10.1021/ol7021998
- Miller, N. A.; Willis, A. C.; Paddon-Row, M. N.; Sherburn, M. S. *Angew. Chem., Int. Ed.* **2007**, *46*, 937–940. doi:10.1002/anie.200603335
- Shimp, H. L.; Hare, A.; McLaughlin, M.; Micalizio, G. C. *Tetrahedron* **2008**, *64*, 3437–3445. doi:10.1016/j.tet.2008.02.015
- Kim, S.; Seomoon, D.; Lee, P. H. *Chem. Commun.* **2009**, 1873–1875. doi:10.1039/b823037d
- Beydoun, K.; Zhang, H.-J.; Sundararaju, B.; Demerseman, B.; Achard, M.; Xi, Z.; Bruneau, C. *Chem. Commun.* **2009**, 6580–6582. doi:10.1039/b913595b
- Englert, M.; Jolly, P. W.; Wilke, G. *Angew. Chem., Int. Ed. Engl.* **1972**, *11*, 136–137. doi:10.1002/anie.197201361
- Pasto, D. J.; Huang, N.-Z. *Organometallics* **1985**, *4*, 1386–1395. doi:10.1021/om00127a014

- Pasto, D. J.; Huang, N.-Z.; Eigenbrot, C. W. *J. Am. Chem. Soc.* **1985**, *107*, 3160–3172. doi:10.1021/ja00297a022
- Arisawa, M.; Sugihara, T.; Yamaguchi, M. *Chem. Commun.* **1998**, 2615–2616. doi:10.1039/a807527a
- Osakada, K.; Choi, J.-C.; Sarai, S.; Koizumi, T.; Yamamoto, T. *Chem. Commun.* **1997**, 1313–1314. doi:10.1039/a701859b
- Lu, P.; Ma, S. *Org. Lett.* **2007**, *9*, 2095–2097. doi:10.1021/ol070632r
- Park, J. H.; Kim, E.; Kim, H.-M.; Choi, S. Y.; Chung, Y. K. *Chem. Commun.* **2008**, 2388–2390. doi:10.1039/b718107h
- Kawamura, T.; Inagaki, F.; Narita, S.; Takahashi, Y.; Hirata, S.; Kitagaki, S.; Mukai, C. *Chem.–Eur. J.* **2010**, *16*, 5173–5183. doi:10.1002/chem.200903568

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
[doi:10.3762/bjoc.7.67](https://doi.org/10.3762/bjoc.7.67)